



Air temperature induced uncertainty in real time flood forecasting over alpine basins

G. Ravazzani^{1*}, A. Ceppi¹, A. Salandin², D. Rabuffetti², M. Mancini¹

¹Politecnico di Milano – D.I.I.A.R., Piazza Leonardo da Vinci 32, 20133, Milano, Italy ²A.R.P.A. Piemonte – Via Pio VII 9, 10135, Torino, Italy *Corresponding author: giovanni.ravazzani@polimi.it



Abstract

Coupling meteorological and hydrological models is recognized by scientific community as a necessary way to forecast extreme hydrological phenomena, in order to active useful mitigation measurements and alert systems in advance.

In order to quantify uncertainty of flood prediction, the hydrological community is increasingly looking at the use of Ensemble Prediction System (EPS) that produce a suite of predictions in contrast to a single forecast of traditional deterministic modelling techniques. Due to an increase in computation power and data transmission rates we are now in a position to use ensemble predictions effectively also for operational flood forecasting, but accurate reliability analysis should be performed.

The goal of this work is to evaluate how the uncertainty of EPS meteorological forecasts influences the performance of hydrological predictions in terms of Quantitative Discharge Forecast (QDF) over alpine basins, focusing the attention on precipitation and air temperature. We show that air temperature is a crucial feature in determining the partitioning of precipitation in solid (snow) and liquid phase (rainfall) and snow melting, therefore having possibility to significantly affect river discharge prediction in autumn and spring seasons even if good accuracy of precipitation forecast was reached.

		-Area	of study w w w w w w w w w w w w w w w w w w w
11 BASINS	Lag time [h]	Area [km ²]	
Ticino basin (closed at Bellinzona)	9	1537	3
Maggia basin (closed at Solduno)	6.8	902	
Toce basin (closed at Candoglia)	9	1534	
Sesia basin (closed at Palestro)	18.8	2606	
Po basin (closed at Carignano)	18	3960	
Stura basin (closed at Fossano)	9.5	1239	Contraction of the second seco
Tanaro basin (closed at Farigliano)	14.8	1457	
Belbo basin (closed at Castelnuovo)	15.1	421	STURA di
Bormida basin (closed at Cassine)	23.2	1523	DEMONTE
Orba basin (closed at Casal Cervelli)	14.2	750	Digital Elevation Model boundaries of each basin
Scrivia basin (closed at Serravalle)	10	617	nauging stations are illust

+6

+8

>+10

The **POLIMI** hydro-meteorological chain:

the forecasting cascade system

Operational real time hydro-meteorological forecast systems are realized by use of one-way coupling, i.e. the meteorological output variables are driven into hydrological models



Hydro-Meteorological data

2000-2008 available database (ARPA Piemonte and Meteo Swiss)

• Temperature: 465 thermometers • Relative Humidity: 186 hygrometers • Precipitation: 486 rain gauge stations • Solar Radiation: 92 pyranometers

• Wind Speed: 123 anemometers

• Hydrometer: 132 data @ basin close sections

Atmospheric forcing

Input values for the FEST-WB model may be derived from observed (i.e. measured) data from the hydro-metrological stations of ARPA Piedmont and of MeteoSwiss or from the forecasted data of the COSMO-LEPS and MOLOCH meteorological models. Afterwards, the FEST-WB simulation calculates different physical processes (snow dynamics, infiltration, water balance, hypodermic and surface propagation etc), and returns different hourly

Meteorological models:

Cosmo-Leps and Moloch

COSMO-LEPS Model (Marsigli et al., 2005)

• Spatial Resolution: 10.0 km (0.09°) • Temporal Resolution: 3 h

• Vertical levels: 40 (non-hydrostatic) • Ensemble members: 16 nested

on ECMWF EPS • Forecast range: +132 h • Run starting at: 12:00 UTC

• Owner: ARPA Emilia-Romagna

MOLOCH Model (Malguzzi et al., 2006)

• Spatial Resolution: 2.3 km (0.02°) • Temporal Resolution: 1 h

• Vertical levels: 50 (non-hydrostatic) • Deterministic model, nested on BOLAM, nested on ECMWF

• Forecast range: +48 h

• Run starting at: 00:00 UTC • Owner: ISAC-CNR



gauging stations are illustrated with vellow dots







ipsographic curve over the Sesia basin even a rising of the temperatures and therefore the snow line, no relevant differences are shown in peak discharges.

Atmospheric Science Letters, 9, 88-94, doi: 10.1002/asl.179, 2008.